



Meriden

ANIMAL HEALTH

Unimectin

1% Injection

Composition

Ivermectin 1.0g
Excipients qsf 100.0ml

Presentation

A sterile non-aqueous solution for injection containing 1% w/v Ivermectin.

Mode of Action

Ivermectin is a broad-spectrum anti-parasite with a wide safety margin. Ivermectin stimulates the release of gamma-aminobutyric acid (GABA), a neurotransmitter from the nerve endings of target parasites. This interrupts the nerve impulses necessary for movement thereby paralyzing and killing the parasites.

Indications

For cattle, sheep and swine Unimectin 1% injection is indicated for the effective treatment and control of the following parasites:

Cattle parasites: *Bunostomum phlebotomum*, *Cooperia spp.*, *Dictyocaulus viviparus*, *Haemonchus placei*, *Nematodirus spp.*, *Oesophagostomum radiatum*, *Ostertagia spp.*, *Thelazia spp.*, *Trichostrongylus spp.*, *Trichuris spp.*, *Hypoderma spp.*, Mange mites, *Psoroptes bovis*, *Sarcoptes scabiei*, Sucking lice, *Linognathus vituli*, *Haematopinus eurysternus*, *Solenopotes capillatus*.

Sheep parasites: *Chabertia ovina*, *Cooperia curticei*, *Dictyocaulus filaria*, *Haemonchus contortus*, *Nematodorus filicollis*, *Oesophagostomum columbianum*, *Ostertagia spp.*, *Protostrongylus rufescerism*, *Trichostrongylus axei*, *T.rufei*, *T.colubriformis*, *T.vitrinus*, *Trichuris ovis*, Nasal bots (*Oestrus ovis*), Mange mites (*Psoroptes ovis*).

Swine Parasites: *Ascaris suum*, *Hyostrongylus rubidus* (adults and forth-stage larvae), *Oesophagostomum spp.* (adults and forth-stage larvae), Threadworm, *Strongyloides ransonii* (adults), *Melaststrongylus spp.* (adults), *Haematopinus suis*, *Sarcoptes scabiei var Suis*.

Dosage & Administration

Unimectin 1% injection should be given only by subcutaneous injection at the recommended dosage level of 200µg Ivermectin per kg body weight under the loose skin in front of, or behind the shoulder in cattle and in the neck in sheep.

Each ml contains 10mg Ivermectin sufficient to treat 50kg body weight of cattle and sheep.

Unimectin 1% injection should be given only by subcutaneous injection in the neck of swine at the recommended dose level of 300µg per kg body weight. Each ml contains 10mg of Ivermectin sufficient to treat 33ka bodv weight of swine.

Storage: Store in a cool and dry place.

Packing: 100ml glass vials

Contra-Indication

This product is not to be used intramuscularly or intravenously

Contact us at:

Meriden Animal Health Limited
Cranfield Innovation Centre
University Way
Cranfield Technology Park
Cranfield MK43 0BT

Tel: +44 (0) 1234 436130
Fax: +44 (0) 1234 436131
Email: sales@meriden-ah.com
Web: www.meriden-ah.com