

DATA SHEET



GESKYPUR®

Presentation

Injectable emulsion. Each 2-ml dose of vaccine contains:

. Aujeszky's disease virus glycoproteins, at least, according to the European Pharmacopoeia	1.1 ΔG7.U
. Benzyl alcohol	10.56 mg
. Oily excipient, q.s.	2 ml

1 ΔG7.U: q.s. to obtain a ΔG7 of 1 during protection against challenge, according to Ph. Eur.

Uses

Inactivated vaccine in adjuvant against Aujeszky's disease (gI-deleted virus subunits).

The vaccine induces an immune status against Aujeszky's disease virus, shown by challenge and by the presence of seroneutralizing antibodies, transmissible by colostral route.

Indication

For breeder pigs and fattening pigs:

Active immunization against Aujeszky's disease.

For piglets:

Passive immunization against Aujeszky's disease by vaccination of pregnant sows.

Dosage and administration

A 2 ml dose injected intramuscularly into the neck in the area behind the ear, according to the following schedule:

In gilts

- Primary vaccination
2 injections with a 21- to 28-day interval.
Perform the second injection at least 8 days before service mating.
- Booster
1 injection 3 to 4 weeks before farrowing.

In young boars

- Primary vaccination
2 injections with a 21- to 28-day interval.

In breeder pigs in activity (sows and boars):

- Boosters
Half-yearly boosters.

In fattening pigs:

- Primary vaccination:
2 injections at the ages of 10 and 14 weeks.

Contra-indications, warnings, etc

There are no contra-indications.

Pharmaceutical precautions

Store between +2°C and +8°C, protected from light.

Further information

Vaccination during pregnancy or lactation is not contraindicated.

No undesirable effect has been observed after the administration of an overdose of vaccine.

Merial Asia Pte Ltd
57A Science Park Drive,
The Faraday
Singapore Science Park 1
Singapore 118238

