

DATA SHEET



HYORESP®

Presentation

Each 2-ml dose of vaccine contains:

. Inactivated whole cell <i>Mycoplasma hyopneumoniae</i> , at least ELISA.U*	3.0
. Al ⁺⁺⁺ (as hydroxide)	4.2 mg
. Thiomersal	0.2 mg
. Excipient, q.s.	2 ml

* ELISA.U: Unit of RP (relative potency) compared to a reference vaccine.

Indication

Immunization of pigs for the reduction of infection and lung lesions of the disease caused by *Mycoplasma hyopneumoniae*.

Dosage and administration

Inject one 2-ml dose of vaccine by intramuscular route, according to one of the following schedules:

- 2 injections at an interval of 3-4 weeks from the age of 5 days.
- 1 injection from the age of ten weeks.

The choice of the vaccination regimen should be based on the knowledge of the risks (time of occurrence of contamination) in the herd. Vaccination must be performed prior to animals' contamination. Thus the 1- injection regimen will be limited to cases where contamination occurs during the mid to late fattening period.

Special Precautions for use

Apply usual procedures for the handling of the animals.

Vaccinate only healthy animals.

Shake before use.

Apply usual aseptic procedures.

Contraindications

No known contraindications

Withdrawal Period

Zero days

Pharmaceutical precautions

Store between +2°C and +8°C, protected from light.

Further information

The injection of the vaccine may cause at the site of injection a transient oedema and muscular inflammation, not usually exceeding one week and seldom up to five weeks. The vaccination may cause a slight temperature increase (less than 1°C) for a transient period (less than 24 hours after injection) without any consequence on the health and the zootechnical performance.

The vaccination may exceptionally reveal a hypersensitivity status. Appropriate treatment for the symptoms shown should be administered.

Special precautions for disposal of unused product and waste material

Since the vaccine is inactivated, no special precaution must be taken concerning unused vaccinal contents.

As all medicinal products, bottles and unused vaccinal contents must be disposed by appropriate channel in compliance with the legislation in force.

Merial Asia Pte Ltd
57A Science Park Drive,
The Faraday
Singapore Science Park 1
Singapore 118238

